

FUNDAMENTAL PLANES, AND THE *BARLESS* $M_{\text{BH}}-\sigma$ RELATION, FOR SUPERMASSIVE BLACK HOLES

ALISTER W. GRAHAM¹

Centre for Astrophysics and Supercomputing, Swinburne University of Technology, Hawthorn, Victoria 3122, Australia.
Draft version February 4, 2008

ABSTRACT

The residuals about the standard $M_{\text{bh}}-\sigma$ relation correlate with the effective radius, absolute magnitude, and Sérsic index of the host bulge. Although, it is noted here that the elliptical galaxies do not partake in such correlations. Moreover, it is revealed that barred galaxies (with their relatively small, faint, and low stellar concentration bulges) can deviate from the $M_{\text{bh}}-\sigma$ relation by $\delta \log M_{\text{bh}} \approx -0.5$ to -1.0 dex (their σ values are too large) and generate much of the aforementioned correlations. Removal of the seven barred galaxies from the Tremaine et al. set of 31 galaxies gives a “barless $M_{\text{bh}}-\sigma$ ” relation with an intrinsic scatter of 0.17 dex (cf. 0.27 dex for the 31 galaxies) and a total scatter of 0.25 dex (cf. 0.34 dex for the 31 galaxies). The introduction of a third parameter does not reduce the scatter. Furthermore, removal of the barred galaxies, or all the disk galaxies, from an expanded and updated set of 40 galaxies with direct black hole mass measurements gives a consistent result, such that $\log(M_{\text{bh}}/M_{\odot}) = (8.25 \pm 0.05) + (3.68 \pm 0.25) \log[\sigma/200 \text{ km s}^{-1}]$. The (barless) $\sigma-L$ relation for galaxies with black hole mass measurements is found to be consistent with that from the SDSS sample of early-type galaxies. In addition the barless $M_{\text{bh}}-\sigma$ relation, the $M_{\text{bh}}-n$ relation, and the $M_{\text{bh}}-L$ relation are shown to yield SMBH masses less than $2\text{--}4 \times 10^9 M_{\odot}$.

Subject headings: black hole physics — galaxies: bulges — galaxies: fundamental parameters — galaxies: structure

1. INTRODUCTION

Tight correlations between supermassive black hole (SMBH) masses and large scale properties of the host bulges are interesting for two obvious reasons. They enable us to predict SMBH masses in thousands of galaxies where the black hole’s sphere-of-influence is highly unresolved, and they provide clues to the physical processes responsible for the co-evolution of black hole and host bulge. Recent endeavors have advocated relations involving not one but two bulge parameters and therein claims of “fundamental planes”, akin to the Fundamental Plane for elliptical galaxies (Djorgovski & Davis 1987; Dressler et al. 1987). The existence of such SMBH fundamental planes imply that current theories for the $M_{\text{bh}}-\sigma$ relation (Ferrarese & Merritt 2000; Gebhardt et al. 2000) or the $M_{\text{bh}}-L$ relation (McLure & Dunlop 2002; Marconi & Hunt 2003; updated in Graham 2007), which do not include a third parameter, are incomplete.

This article investigates the fundamental planes for SMBHs² involving the parameters M_{bh} , R_{e} , and $\langle \mu \rangle_{\text{e}}$ (Barway & Kembhavi 2007), M_{bh} , R_{e} , and σ (Marconi & Hunt 2003; de Francesco et al. 2006; Aller & Richstone 2007; Hopkins et al. 2007), and, for the first time, M_{bh} , σ , and n . In Section 2 it is explained why a previous claim for a small ‘total’ scatter (0.19 dex) about the $M_{\text{bh}}-R_{\text{e}}-\langle \mu \rangle_{\text{e}}$ plane was the result of a miscalculation. In Section 3 it is revealed that the galaxies which deviate from the $M_{\text{bh}}-\sigma$ relation, giving rise to the $M_{\text{bh}}-\sigma-R_{\text{e}}$, $M_{\text{bh}}-\sigma-L$, and $M_{\text{bh}}-\sigma-n$ relations with less scatter than the $M_{\text{bh}}-\sigma$ relation, are predominantly barred galaxies. A “barless $M_{\text{bh}}-\sigma$ ” relation, and an elliptical-only $M_{\text{bh}}-\sigma$ relation,

is subsequently constructed in Section 4 and found to heavily nullify the evidence for fundamental planes for SMBHs and their host bulges.

Given the recent discussion in the literature about biases in the $M_{\text{bh}}-\sigma$ and/or $M_{\text{bh}}-L$ relation, and also in the local sample of galaxies with direct SMBH mass measurements, these concerns are explored here. In Sections 5 a $\sigma-L$ relation is constructed and shown to be equal to that obtained using SDSS early-type galaxy data, thereby laying to rest concerns that the local sample of galaxies with direct SMBH mass measurements may be biased with respect to the greater population (e.g. Yu & Tremaine 2002; Bernardi et al. 2007). Furthermore, in section 6, the K -band $M_{\text{bh}}-L$ relation and the barless $M_{\text{bh}}-\sigma$ relation are shown to yield consistent results with neither giving SMBH masses greater than $\sim 4 \times 10^9 M_{\odot}$.

2. THE $(M_{\text{BH}}, \langle \mu \rangle_{\text{E}}, R_{\text{E}})$ PLANE

Barway & Kembhavi (2007, hereafter BK07) made the interesting claim that a combination of two photometric parameters, namely the effective radius R_{e} and the mean effective surface brightness $\langle \mu \rangle_{\text{e}} = -2.5 \log \langle I \rangle_{\text{e}}$, can be used to predict SMBH masses with a greater degree of accuracy than single quantities such as luminosity or velocity dispersion.

A tight relation exists between black hole mass, M_{bh} , and the luminosity, L , of the host bulge. The luminosity can of course be expressed in terms of two other parameters because $L = 2\pi R_{\text{e}}^2 \langle I \rangle_{\text{e}}$, where $\langle I \rangle_{\text{e}}$ is average intensity within the effective half light radius, R_{e} , of the bulge. One question of interest is whether the scatter about the $M_{\text{bh}}-(R_{\text{e}}^2 \langle I \rangle_{\text{e}})$ relation can be reduced by allowing the exponents on $\langle I \rangle_{\text{e}}$ and R_{e} to deviate from their 1:2 ratio and, importantly, if this results in less scatter than the other competing relations. Given the small scatter about the Fundamental Plane — involving R_{e} ,

¹ Corresponding Author: AGraham@astro.swin.edu.au

² The “fundamental plane of black hole activity” involving radio core luminosity and X-ray luminosity (Merloni et al. 2003; Falcke et al. 2004) is not addressed here.

$\langle\mu\rangle_e = -2.5 \log\langle I\rangle_e$, and σ — and the tight relationship between M_{bh} and σ (Merritt & Ferrarese 2001; Tremaine et al. 2002), one may indeed expect a well defined plane using the parameters R_e , $\langle\mu\rangle_e$, and M_{bh} .

2.1. Barway & Kembhavi data

This section examines BK07’s claim that the total root mean square (r.m.s.) scatter in the $\log M_{\text{bh}}$ direction, when using $\langle\mu\rangle_e$ and $\log R_e$ as predictor quantities of M_{bh} , is 0.25 dex (and 0.19 dex when excluding the outlier NGC 4742).

A simple linear, $Y = A + BX$, ordinary least squares regression analysis $\text{OLS}(Y|X)$ is performed with $Y = \log M_{\text{bh}}$ and $X = \log R_e + b\langle\mu\rangle_e$. Solving for the parameters A , B , and b , this non-symmetrical regression gives the smallest r.m.s. residual in the $\log M_{\text{bh}}$ direction, which is what one wants when using the $M_{\text{bh}}-\langle\mu\rangle_e-R_e$ plane to predict M_{bh} in other galaxies (see Feigelson & Babu 1992). The data for $\log M_{\text{bh}}$, $\log R_e$, and $\langle\mu\rangle_e$ have been taken from Table 1 in BK07. Due to the absence of reported errors on the quantities $\log R_e$ and $\langle\mu\rangle_e$ in BK07, measurement errors are not included in the regression, and subsequently no attempt to quantify the intrinsic scatter has been made. Parameter uncertainties are derived here using a bootstrap sampling of the data points (i.e. sampling with replacement from the original sample) to produce 1000 Monte Carlo samples from which 1000 optimal fits are derived. This provides a histogram of each parameter from which one can compute the central 68.3% width, which is used as the 1σ uncertainty.

The optimal (B -band) solution using all 18 data points from BK07 is

$$\log(M_{\text{bh}}/M_\odot) = (8.18 \pm 0.09) + (3.15 \pm 0.33) \log[R_e/3 \text{ kpc}] - (0.90 \pm 0.18)[\langle\mu\rangle_{e,B} - 21.0]. \quad (1)$$

The total scatter in the $\log M_{\text{bh}}$ direction is 0.32 dex. However the total scatter in the $M_{\text{bh}}-\sigma$ relation for this same galaxy set is 0.31 dex.

BK07 performed an additional analysis, excluding NGC 4742 whose SMBH mass derivation has not yet appeared in a refereed paper (see Tremaine et al. 2002) and may therefore potentially be erroneous. This galaxy also appeared as a clear outlier in their data. This does not necessarily mean the data point is in error; it may simply be a 3σ event, or the distribution of residuals may perhaps not be ‘normal’. Robust statistics requires that outlying data points not bias an analysis. No single data point from a distribution should have the ability to significantly alter the result of an analysis dictated by the remaining population. The optimal relation after excluding NGC 4742 is given by the expression

$$\log(M_{\text{bh}}/M_\odot) = (8.21 \pm 0.07) + (3.23 \pm 0.26)[\log R_e/3 \text{ kpc}] - (1.01 \pm 0.13)[\langle\mu\rangle_{e,B} - 21.0] \quad (2)$$

with a total scatter of 0.25 dex. The low value of 0.19 dex reported by BK07 appears to have arisen by dividing their scatter in the $\log R_e$ direction (0.061) by the coefficient in front of the $\log M_{\text{bh}}$ term in their eq.3 (which is their fitted plane). However, this approach overlooks the three-dimensional nature of the plane and consequently results in an over-estimation of the plane’s ability to predict black hole masses. Computing the r.m.s. offset between the black hole masses listed in Table 1 of

BK07 (excluding NGC 4742) and the values predicted from their plane (their eq.3), which can be re-written as $\log(M_{\text{bh}}/M_\odot) = 8.28 + 3.13[\log R_e/3 \text{ kpc}] - 0.97[\langle\mu\rangle_{e,B} - 21]$, one obtains a total scatter in the $\log M_{\text{bh}}$ direction of 0.27 dex, not 0.19 dex, and greater than the value of 0.25 dex obtained above using Eq.2.

Although the claim in BK07 appears misplaced, based on an erroneous treatment of the data, the idea tested there is a valid one. In an effort to improve the $M_{\text{bh}}-\langle\mu\rangle_e-R_e$ plane’s reliability for predicting black hole masses, it is noted here that three of the galaxies used by BK07 are known disk galaxies, or at least they are not regular elliptical galaxies. M32 may be a stripped S0 galaxy (Bekki et al. 2001; Graham 2002), while NGC 2778 is a disk galaxy (Rix, Carollo & Freeman 1999) as is NGC 4564 (Trujillo et al. 2004; see also figure 6 in Graham & Driver 2007a). Consequently, the effective radii and mean surface brightnesses which have been used for these three galaxies do not pertain to their bulges. IC 1459 is also excluded here due to the order of magnitude uncertainty on its SMBH mass (Cappellari et al. 2002). Excluding these four galaxies plus NGC 4742 gives, from a reduced sample of only 13 galaxies, a total scatter of 0.28 dex. However, the total scatter in the $M_{\text{bh}}-\sigma$ relation for this cleaned galaxy set is 0.27 dex. The $M_{\text{bh}}-\sigma$ relation therefore appears more competitive than the $M_{\text{bh}}-\langle\mu\rangle_e-R_e$ plane.

This sample size is obviously small and therefore makes it hard to reach reliable conclusions. The $M_{\text{bh}}-\langle\mu\rangle_e-R_e$ plane is thus investigated further with a larger galaxy sample in the following subsection.

2.2. Marconi & Hunt data

Instead of using the B -band data in BK07, the scatter about the $M_{\text{bh}}-\langle\mu\rangle_e-\log R_e$ plane is explored here using the larger, homogeneous, K -band data set from Marconi & Hunt’s (2003; hereafter MH03) 27 “Group 1” galaxies. Using the minor-to-major axis ratio, b/a , of the GALFITted (Peng et al. 2002) Sérsic bulge component (Marconi & Hunt, priv. comm.), MH03’s tabulated major-axis effective radii, $R_{e,\text{maj}}$, have been converted into a geometric mean radius $R_e = \sqrt{R_{e,\text{maj}}^2(b/a)}$ which is used here. While MH03 did not report any values for $\langle\mu\rangle_e$, they can be derived from the expression

$$\langle\mu\rangle_e = m_{\text{tot}} + 2.5 \log[2\pi R_{e,\text{maj}}^2(b/a)], \quad (3)$$

where m_{tot} is the apparent magnitude of the bulge (obtained from the absolute magnitude and distance in Table 1 in MH03). These values are shown in Table 1.

Four of the five galaxies which were excluded at the end of Section 2.1 are in MH03’s “Group 1” list. They are again excluded here for the same reasons. Doing so, one obtains from the remaining 23 galaxies, using the M_{bh} values given in MH03,

$$\log(M_{\text{bh}}/M_\odot) = (7.92 \pm 0.12) + (2.24 \pm 0.37)[\log R_e/3 \text{ kpc}] - (0.54 \pm 0.14)[\langle\mu\rangle_{e,K} - 17.5], \quad (4)$$

which has a total scatter in the $\log M_{\text{bh}}$ direction of 0.33 dex. The coefficients in Eq.4 are consistent with an $M_{\text{bh}}-(I_e R_e^2)$ plane, and the total scatter in the $M_{\text{bh}}-L$ relation for these galaxies (0.34 dex) is comparable. Moreover, the total scatter in the $M_{\text{bh}}-\sigma-R_e$ plane for this same

galaxy set is 0.28 dex. It is therefore concluded that the $M_{\text{bh}}-\langle\mu\rangle_e-\log R_e$ plane is not warranted.

The following section explores the $M_{\text{bh}}-\sigma-R_e$ plane, and other planes involving M_{bh} , σ and some third parameter.

3. THE $(M_{\text{BH}}-\sigma)-X$ PLANE

MH03 explored the addition of $\log R_e$ to the $M_{\text{bh}}-\sigma$ relation to create a “fundamental plane for SMBHs”. From their 27 “Group 1” galaxies, they constructed a relation between M_{bh} and $R_e\sigma^2$ (proportional to the virial bulge mass), which resulted in an intrinsic dispersion³ (total scatter) of 0.25 (0.30) dex in the $\log M_{\text{bh}}$ direction. Allowing the exponents on the R_e and σ terms to vary independently, Hopkins et al. (2007) used the same 27 Group 1 galaxies from MH03 along with some updated measurements, to report that $\log(M_{\text{bh}}/M_\odot) = (8.33 \pm 0.06) + (0.43 \pm 0.19) \log[R_e/3 \text{ kpc}] + (3.00 \pm 0.30) \log[\sigma/200 \text{ km s}^{-1}]$, with an intrinsic scatter of 0.21 dex (and a total scatter of 0.30 dex, Hopkins 2007, priv. comm.). For comparison, the $M_{\text{bh}}-\sigma$ relation in Tremaine et al. (2002) has an intrinsic (total) scatter of 0.27 (0.34) dex. It therefore appears that the introduction of a third parameter to the standard $M_{\text{bh}}-\sigma$ relation may reduce the scatter and Hopkins et al. (2007) show that it does. Here it is investigated which third parameter is optimal.

From Graham & Driver (2007a, hereafter GD07) the total scatter about the log-quadratic M-n relation is reported to be 0.31 dex. This highlights the strong connection between M_{bh} and the radial structure in the stellar distribution of the host bulge (see also Graham et al. 2007, their Section 1), and hence the need to advance beyond $R^{1/4}$ models and their associated luminosity/mass dependent biases (e.g., Trujillo et al. 2001; Brown et al. 2003). This Section explores whether the scatter about the $M_{\text{bh}}-\sigma$ relation is best reduced through the addition of the host bulge’s $\log R_e$, K -band magnitude, or $\log n$. The largest homogeneous sample of galaxies with published M_{bh} and n values is that in GD07. One of the strengths of the $M_{\text{bh}}-n$ relation is that photometrically uncalibrated images can be used. While this means that GD07 do not have magnitudes, they do have bulge R_e values and bulge-to-total ratios which can, when needed, be applied to the galaxy M_K values in MH03. The Sérsic indices from GD07 pertain to the major-axis. It is perhaps worth noting that from very early on it was known that the major- and minor-axis need not and do not have the same Sérsic profile shape (Caon et al. 1993). In the presence of ellipticity gradients the Sérsic index will vary with position angle (Ferrari et al. 2004), and the value obtained from a symmetrical 2D fit with a single Sérsic index will match neither the major- nor minor-axis value. Moreover, the random viewing angles at which spheroids are viewed will also introduce scatter to the $M_{\text{bh}}-n$ relation (and the $M_{\text{bh}}-\sigma$ relation if the bulges are triaxial).

The Sérsic indices and SMBH masses for the 27 galaxies tabulated in GD07 are used here, along with the (geometric mean) effective radii (Table 1), K -band magnitudes from MH03, and the central velocity disper-

sions from Ferrarese & Ford (2005, hereafter FF05). NGC 6251 and NGC 7052 had a different distance in GD07 and MH03, and have had their R_e and M_K values adjusted to match the distances in GD07. Although MH03 did not include/model NGC 1399 (Houghton et al. 2006), a velocity dispersion $\sigma = 344 \text{ km s}^{-1}$ (HyperLeda) has been adopted, along with the Sérsic index, R_e value, and B -band magnitude from D’Onofrio et al. (1994), adjusted to a distance of 20 Mpc (Tonry et al. 2001), and using $b/a = 0.94$ (NED) and $B - K = 4.14$ (Buzzoni 2005). The bulge parameters from Graham (2002) are used for NGC 221, along with a Johnson $R - K$ color of 2.34 (Buzzoni 2005). This left two galaxies (NGC 2778 and NGC 4564) which had R_e and M_K values pertaining to the galaxy rather than the bulge in MH03. From the analysis in GD07, the values $R_{e,\text{maj}}$ equals 0.25 and 0.31 kpc, and the B/T ratios 0.21 and 0.24 have been adopted, respectively.

For these 27 galaxies one obtains an $M_{\text{bh}}-\sigma$ relation similar to that reported in Tremaine et al. (2002); it is such that

$$\log(M_{\text{bh}}/M_\odot) = (8.09 \pm 0.07) + (4.08 \pm 0.40) \log[\sigma/200 \text{ km s}^{-1}], \quad (5)$$

with Δ , the total r.m.s. scatter in the M_{bh} direction, equal to 0.31 dex.

The $M_{\text{bh}}-\sigma-L$ plane for these galaxies is given by

$$\log(M_{\text{bh}}/M_\odot) = (8.13 \pm 0.07) - (0.11 \pm 0.05)[M_K + 24] + (3.34 \pm 0.48) \log[\sigma/200 \text{ km s}^{-1}], \Delta = 0.27 \text{ dex} \quad (6)$$

The $M_{\text{bh}}-\sigma-R_e$ plane is

$$\log(M_{\text{bh}}/M_\odot) = (8.15 \pm 0.06) + (0.28 \pm 0.12) \log[R_e/3 \text{ kpc}] + (3.65 \pm 0.32) \log[\sigma/200 \text{ km s}^{-1}], \Delta = 0.26 \text{ dex} \quad (7)$$

while the $M_{\text{bh}}-\sigma-n$ plane is

$$\log(M_{\text{bh}}/M_\odot) = (7.98 \pm 0.05) + (1.11 \pm 0.32) \log[n/3] + (2.72 \pm 0.52) \log[\sigma/200 \text{ km s}^{-1}], \Delta = 0.23 \text{ dex} \quad (8)$$

Given the best performer is the $M_{\text{bh}}-\sigma-n$ plane, the residuals about the $M_{\text{bh}}-\sigma$ relation are plotted in Fig.1 against the bulge Sérsic index. From Fig.1a it is clear that a trend will still persist after the exclusion of the five galaxies whose SMBH sphere-of-influence is not resolved (according to Table II from FF05). On the other hand, Fig.1b reveals that the trend is caused by (some of) the disk galaxies. This intriguing aspect is explored further in the following section.

4. THE “BARLESS $M_{\text{BH}}-\sigma$ ” RELATION

4.1. Graham & Driver data

Looking at Fig.1, much of the trend is due to some five data points from the small bulges of disk galaxies. While these five bulges have small ($R_e < 1 \text{ kpc}$) effective radii, some of the other disk galaxies have comparable radii but do not deviate from the $M_{\text{bh}}-\sigma$ relation. These five systems have SMBH masses ~ 0.5 dex below the best fitting $M_{\text{bh}}-\sigma$ relation. Intriguingly, all of these five disk galaxies have been identified in the literature as containing bars. They are: the Milky Way (e.g., López-Corredoira et al. 2007, their Fig.5); NGC 1023 (Debattista et al. 2002, their Fig. 4b; Sil’chenko 1999); NGC 2778 (weak bar peaks at $5''$; Rest et al. 2001, their Fig. 8; Trujillo et

³ Intrinsic dispersion, sometimes called internal dispersion, is the scatter remaining after subtracting in quadrature, from the total scatter, the contribution from the assumed measurement uncertainties.

al. 2004); NGC 2787 (Erwin et al. 2003, their Fig. 1); and NGC 3384 (Busarello et al. 1996, their Fig. 7; Cappellari & Emsellem 2004; Erwin 2004; Meusinger & Ismail 2007). In sharp contrast to this, only one of the other 8 disk galaxies (NGC 4258, van Albada 1980) is classified in NED as having a bar; therefore, if any of the other seven disk galaxies do possess a bar, it must be weak. If the probability of a disk galaxy having a bar is equal to the probability of not having a bar, then the distribution in Figure 1 has a 1 in 1024 chance of occurring. If 75% of disk galaxies have bars (e.g., Eskridge et al. 2000; see also Knapen et al. 2000 and Marinova & Jogee 2007), then the observed distribution has less than a one in ten thousand likelihood of occurring by chance.

In Fig. 2, the barred galaxies can be seen to be largely responsible for the reduced scatter when going from the $M_{\text{bh}}-\sigma$ relation to the $M_{\text{bh}}-\sigma-R_e$, $M_{\text{bh}}-\sigma-L$, and $M_{\text{bh}}-\sigma-n$ planes. In other words, these galaxies deviate from the $M_{\text{bh}}-\sigma$ relation. It is therefore of interest to re-derive the $M_{\text{bh}}-\sigma$ relation excluding those galaxies with bars. For the 21 non-barred galaxies in GD07

$$\log(M_{\text{bh}}/M_{\odot}) = (8.20 \pm 0.05) + (3.83 \pm 0.10) \log[\sigma/200 \text{ km s}^{-1}], \quad (9)$$

with a total scatter of 0.22 dex (cf. Eq. 5). Aside from NGC 4258, the barred galaxies have an offset in the $\log M_{\text{bh}}$ direction of 0.5 to 0.8 dex.

In passing it is noted that a prolate bulge, even in the absence of a bar, will have a smaller effective radius when viewed along its major-axis (e.g., Lanzoni & Ciotti 2003). A more detailed investigation of the above galaxies could therefore include how the measured size, magnitude, and concentration of the spheroidal component, and the velocity dispersion, changes with the orientation of the bulge and bar.

Although Figures 2 might appear to hint that the barred galaxies have smaller effective radii than the non-barred disk galaxies, a KS test reveals no significance (at even the 1σ level) that the cumulative distribution function for the barred and unbarred disk galaxies' effective radii may be different. Similarly, Student's t-test reveals no significant difference between the means of each distribution, with only an 86% ($<1.5\sigma$) probability of difference.

It is pertinent to ask whether the inclusion of an additional parameter to the above “barless $M_{\text{bh}}-\sigma$ ” relation is warranted. The answer appears to be ‘no’. Reductions of not more than 0.01 dex are achieved through the addition of either R_e , L , or n . This implies that a fundamental plane for SMBHs is not appropriate; if it was, it should equally apply to galaxies with and without bars.

A bar may result in the fueling of the SMBH (e.g., Wyse 2004; Ohta et al. 2007), perhaps eventually bringing its mass in line with the $M_{\text{bh}}-\sigma$ relation. Although, the large ratio of barred galaxies to active galaxies in the Universe today would seem to argue against this as a common phenomenon, as does the incidence of bars in Seyfert and normal disk galaxies (Mulchaey & Regan 1997; Ho et al. 1997; although see Crenshaw et al. 2003). It may however be that some other physical property such as nuclear disks or kinematically decoupled cores are influencing the measured velocity dispersions.

Bar instabilities are believed to lead to the formation of pseudobulges. Such evolution may have resulted in

(pseudo)bulges with an increased velocity dispersion and luminosity but a relatively anaemic SMBH (unless it also grew during the formation of the pseudobulge). If the barred galaxies do indeed have discrepantly low SMBH masses, rather than high σ values, they should also appear as systematic outliers in the $M_{\text{bh}}-L$ diagram. Figure 3 reveals that this is not the case, suggesting that the SMBH masses are okay but the velocity dispersions are discrepant.

Perhaps the bar dynamics have biased the measurement of the bulge velocity dispersion. The non-circular (streaming) motions of stars in bars obviously deviates from that of the random motions of the bulge stars and may potentially interfere with the central velocity dispersion measurements. An alignment of the (radial orbits in the) bar with our line-of-sight may result in such a scenario⁴. Indeed, the barred galaxy NGC 3384 is highly inclined and has the bar closely aligned with the projected minor axis (Busarello et al. 1996; Erwin et al. 2004). In the case of NGC 1023, a quick visual inspection reveals a disk/bar position angle of $80^\circ/72^\circ$, suggesting that the inclination of this galaxy does not result in us looking down the length of the bar. However, as Debattista (2002) revealed, after deprojecting this galaxy, it was found to have a strong bar whose position angle is 102° offset from the galaxy's (projected) major-axis. That is, we *are* in fact looking down the barrel of the bar in this galaxy. A full treatment of each of the barred galaxies is beyond the scope of this paper, it is however noted that the (projected) bar position angles can appear more aligned with the (projected) major-axis than they are in reality, as is the case with NGC 1023.

4.2. Tremaine et al. data

Removing the seven barred galaxies (the six mentioned above plus NGC 4596) from the Tremaine et al. (2002) sample of 31 galaxies (with the SMBH mass for NGC 821 updated with the value in Richstone et al. 2007), application of Tremaine et al.'s regression technique gives

$$\log(M_{\text{bh}}/M_{\odot}) = (8.21 \pm 0.05) + (3.89 \pm 0.26) \log[\sigma/200 \text{ km s}^{-1}], \quad (10)$$

with an intrinsic scatter of 0.17 dex (cf. 0.27 dex using the original 31 galaxies) and a total scatter of 0.25 dex (cf. 0.34 dex using the original 31 galaxies). This is in agreement with Eq. 9 which used a slightly different sample and velocity dispersions from FF05.

Construction of a barless $M_{\text{bh}}-\sigma-M_B$ plane, with the absolute B -band magnitude, M_B , taken from Tremaine et al. (2002), has the same scatter as the barless $M_{\text{bh}}-\sigma$ relation. Furthermore, the intrinsic scatter from Eq. 10 is smaller than the value of 0.21 dex reported in Hopkins et al. (2007). It seems reasonable to conclude that previous claims for the existence of “fundamental planes for SMBHs” have been influenced by the presence of barred galaxies.

Using the Tremaine et al. (2002) sample and performing a regression which minimizes the residuals in the $\log \sigma$ direction, rather than the $\log M_{\text{bh}}$ direction, the intercept and slope of the barless $M_{\text{bh}}-\sigma$ relation are 8.21 and 4.05. A symmetrical regression will therefore have a

⁴ Velocity dispersion drops (Márquez et al. 2003; Wozniak et al. 2003) would, however, work in the opposite sense.

slope around $(3.89+4.05)/2 = 3.97$. Using the symmetrical bisector linear regression routine BCES from Akritas & Bershady (1996), one obtains

$$\log(M_{\text{bh}}/M_{\odot}) = (8.20 \pm 0.05) + (3.95 \pm 0.25) \log[\sigma/200 \text{ km s}^{-1}]. \quad (11)$$

This expression should be preferred when trying to understand the physical processes responsible for the correlation (see Feigelson & Babu 1992), although this point is perhaps moot given Eq.11's consistency with Eq.9 and 10.

4.3. Marconi & Hunt data

The analysis in Hopkins et al. used the 27 ‘Group 1’ galaxies from MH03. In an effort to better understand the result in Hopkins et al., the original data from MH03 is analyzed here. Given that MH03's M_{bh} and σ values for the Milky Way and M31 were not in dispute (only their K -band magnitudes were somewhat in doubt), these two galaxies have been included here with the 27 ‘Group 1’ galaxies.

The residuals, in the $\log M_{\text{bh}}$ direction, about the $M_{\text{bh}}-\sigma$ relation are shown in Figure 4a for the above 29 galaxies. The trend between the residuals and the effective radii of the host spheroids does indeed appear to suggest the need for a fundamental plane type relation, akin to that proposed by MH03, and later by Hopkins et al. (2007) and also Aller & Richstone (2007) using a sample of 23 galaxies. However, once one identifies the (five) barred galaxies in the above sample, the evidence for such a plane is reduced. The three galaxies with the largest negative residual are barred galaxies.

The three galaxies with the highest positive residuals in Figure 4a — two of which still seem to advocate the need for a ‘fundamental plane’ — are, in order of increasing R_e : the radio galaxy Centaurus A, the Seyfert galaxy NGC 5252 at ~ 100 Mpc, and Cygnus A at a distance of 240 Mpc. The SMBH mass estimate for Centaurus A that was used by MH03 and used in Figure 4 has however since been revised downward by more than a factor of two (Marconi et al. 2006) and so Cen A is therefore now known not to be an outlier. Due to their distances⁵ and somewhat disturbed morphology, none of these galaxies had been fitted with a Sérsic profile by GD07, nor had they been included in Tremaine et al. (2002). While one can conclude that (some) barred galaxies deviate significantly from the $M_{\text{bh}}-\sigma$ relation, the inclusion of NGC 5252 and Cygnus A may present some evidence in favor of a fundamental plane for SMBHs.

If a fundamental plane for black holes does exist, demonstrating its existence with an elliptical only sample would help eliminate concerns that unrelated processes pertaining to bars and disks are misleading us. Figure 4b shows the residuals about the $M_{\text{bh}}-\sigma$ relation for the 17 elliptical galaxies from the sample of 29. One can immediately see that there is, as yet, no convincing evidence for an elliptical galaxy SMBH fundamental plane involving M_{bh} , σ , and R_e . Given the obvious need for more data, the following section introduces and uses new SMBH data obtained after 2003.

4.4. Additional data

Since MH03's paper, additional galaxies have had their SMBH masses measured. These are provided in Table 2, along with galaxies from MH03 for which some updates have become available, giving a total of 40 galaxies with direct SMBH mass measurements. An additional 15 galaxies with somewhat uncertain SMBH mass estimates (see FF05) are listed in Table 3. Although these are not used here, they are provided for a sense of awareness as to further galaxies which may be useful in the future.

Using (i) the (updated) data for the 27 ‘Group 1’ galaxies from MH03 (except for IC 1459 and NGC 4594, whose M_{bh} values are somewhat uncertain), plus (ii) MH03's ten ‘Group 2’ galaxies (minus NGC 1068, NGC 4459, and NGC 4596 for which the SMBH mass estimates are also not secure), plus (iii) the nine new galaxies in Table 2 (excluding NGC 2748 for which there is no published velocity dispersion), gives a total sample of $(25+7+8=)$ 40 galaxies from which an updated $M_{\text{bh}}-\sigma$ diagram has been constructed (Figure 5). For these galaxies, the $M_{\text{bh}}-\sigma$ relation and the $M_{\text{bh}}-\sigma-R_e$ plane are given in Table 4, along with the associated total scatter. For the full data set, one obtains an $M_{\text{bh}}-\sigma$ relation in good agreement with Tremaine et al. (2002). One also has $M_{\text{bh}} \propto \sigma^{3.23 \pm 0.28} R_e^{0.43 \pm 0.11}$, in agreement with the result in Hopkins et al. (2007).

However, from Fig. 5 one can clearly see that many of the barred galaxies deviate from the $M_{\text{bh}}-\sigma$ relation and are obviously responsible for some of the perceived need for a fundamental plane. If these galaxies had R_e values that were smaller than any of the other bulges, then one could argue that the presence of the bar may have nothing to do with their displacement from the $M_{\text{bh}}-\sigma$ relation, and that a ‘fundamental plane’ is needed. However this is not the case, that is, other small spheroids exist which do not deviate from the $M_{\text{bh}}-\sigma$ relation. The only non-barred galaxy with a notable negative $\delta \log M_{\text{bh}}$ residual in Figure 5 and 6 is the LINER galaxy NGC 3998 (De Francesco et al. 2006). As remarked by Fisher (1997), this galaxy has a very steep central velocity dispersion profile, dropping from $\sim 320 \text{ km s}^{-1}$ at $r = 0$ to $\sim 160 \text{ km s}^{-1}$ at $r = 4$ arcseconds (270 pc). A velocity dispersion of 210 (or 250) km s^{-1} for this galaxy would result in a zero (or 1σ) residual about the $M_{\text{bh}}-\sigma$ relation.

Removing the 11 barred galaxies from the sample of 40, one obtains the ‘barless’ $M_{\text{bh}}-\sigma$ relation given in Table 4. The vertical residuals about this relation are shown in Figure 6a, along with the offsets of the barred galaxies relative to this $M_{\text{bh}}-\sigma$ relation defined by the non-barred galaxies. While seven of the ten barred galaxies with known R_e values are responsible for much of the trend between the $M_{\text{bh}}-\sigma$ residuals and R_e , the non-barred galaxies do still reveal a trend. Indeed, from Table 4, one can see that the non-barred galaxies favor a fundamental plane relation. However, it is noted that removal of just two galaxies (NGC 3998 and Cygnus A) from the sample of 29 non-barred galaxies leaves the coefficient in front of the $\log(R_e/3)$ term inconsistent with a value of zero at a significance of less than 2σ . It is disconcerting that just a couple of points are responsible for the apparent plane. The bulk of the data does not suggest the need for a fundamental plane.

As noted previously, to be certain that a 3-parameter fundamental plane is required to describe the connection between SMBHs and their host spheroids, one would ide-

⁵ The third and only other galaxy further than 35 Mpc is NGC 6251 at 107 Mpc.

ally like to use a sample of elliptical galaxies. This would ensure that the ‘plane’ is not a byproduct of additional physical mechanisms or biases related to the presence of a disk and/or bar. Using the 19 elliptical galaxies from the sample of 40 galaxies, one has

$$\log(M_{\text{bh}}/M_{\odot}) = (8.25 \pm 0.05) + (3.68 \pm 0.25) \log[\sigma/200 \text{ km s}^{-1}], \quad (12)$$

with a total scatter of 0.24 dex. Exclusion of the single data point for Cygnus A, the galaxy with the greatest residual offset in Fig.6b, reduces the total scatter to 0.18 dex. This is the same scatter as that about the best fitting $M_{\text{bh}}-\sigma-R_e$ plane to this set of 18 elliptical galaxies. The elliptical galaxies therefore do not provide substantial support for the existence of an $M_{\text{bh}}-\sigma-R_e$ fundamental plane for SMBHs (see Table 4). When using all 19 elliptical galaxies, the 2σ uncertainty on the coefficient in front of the R_e term ranges from -0.08 to 0.55. This parameter is inconsistent with a value of zero at only the 1.4σ level. Moreover, removing just one data point (Cygnus A) reduces the coefficient in front of the R_e term to 0.09 ± 0.11 .

Given the small sample sizes involved, it may be premature to completely rule out the existence of a fundamental plane for SMBHs. Some may object to the removal of outlying data points, which is why equations using both complete and adjusted data sets have been provided. Most would however acknowledge that a certain degree of caution must be associated with any conclusion that hinges on outlying data points. Indeed, for similar reasons, 3σ clipping of distributions is a somewhat common practice these days. One thing which is clear is that the biasing presence of disc (especially barred) galaxies appear responsible for much of the alleged evidence for requiring a fundamental plane for SMBHs.

One should not use either the “barless $M_{\text{bh}}-\sigma$ ” relation nor the standard $M_{\text{bh}}-\sigma$ relation for barred galaxies because the resultant SMBH mass estimates may be in error (too high) by 0.5 to 1.0 dex. One should also not apply an $M_{\text{bh}}-\sigma-R_e$ fundamental plane in the hope of accounting for barred galaxies because such a plane will introduce a bias to the non-barred galaxies.

5. THE σ - L RELATION AND SAMPLE BIAS

There has been some concern recently that the $M_{\text{bh}}-\sigma$ and/or $M_{\text{bh}}-L$ relation may be biased, and that they are not consistent with each other. Lauer et al. (2007), Bernardi et al. (2007), and Graham (2007, his Appendix A) have reported a slight difference in the σ - L relation between the local sample of galaxies with direct SMBH masses and the greater population. If correct, this implies that either the $M_{\text{bh}}-\sigma$ or $M_{\text{bh}}-L$ relation may be biased. Given the offset nature of some of the barred galaxies in the $M_{\text{bh}}-\sigma$ diagram — in the sense that they have overly large velocity dispersions for their SMBH masses — it is apposite to explore if the barred galaxies may be responsible for the allegedly biased nature of these local inactive galaxy samples.

The Group 1 and 2 galaxy data from MH03 is used here, along with the updates noted in Table 2. The 7 barred galaxies from MH03’s 37 ‘Group 1’ and ‘Group 2’ galaxies are excluded, and the K -band magnitudes have been converted to the R_c -band using $R_c - K = 2.6$ (Buzoni 2005). An uncertainty of 0.3 mag and 5% is assigned

to the magnitudes and velocity dispersions, respectively. Applying the regression analysis scheme from Tremaine et al. (2002) to minimize the scatter in the $\log \sigma$ direction, the optimal σ - L relation is

$$\log \sigma = 2.23 \pm 0.03 - (0.092^{+0.018}_{-0.012})[M_R + 21], \quad (13)$$

which is shown in Figure 7. The parameter uncertainties have been estimated from a Monte Carlo bootstrap analysis. Although MH03 note that the M_K value for M31 may be in error, excluding it from the regression has no effect on Eq.13. However, the extreme outlying point NGC 4342, the smallest and faintest spheroid from MH03’s sample after M32, is excluded from this regression.

The reason for constructing an R_c -band relation was to allow a comparison with the result from Tundo et al. (2007, their Eq.4), which is a SDSS r' -band σ - L relation for early-type SDSS galaxies, the majority of which presumably do not have bars. Using $r' - R_c = 0.24$ (Fukugita et al. 1995), Tundo et al.’s expression is such that $\log \sigma = 0.27 - 0.092 M_{R_c} = 2.20 - 0.092(M_{R_c} + 21)$, in remarkable agreement with Eq.13. Therefore, it is not yet established that the local sample of galaxies with direct SMBH mass measurements is biased.

6. $M_{\text{BH}}-\sigma$ VERSUS $M_{\text{BH}}-L$

Given that the local (predominantly inactive) sample of galaxies with direct SMBH mass measurements appears to be unbiased with respect to the greater population, it is appropriate to re-examine whether the (barless) $M_{\text{bh}}-\sigma$ and $M_{\text{bh}}-L$ relations predict different SMBH masses. Indeed, it has been claimed that these relations are not consistent with each other, in the sense that massive galaxies are predicted to have more massive SMBHs when using the $M_{\text{bh}}-L$ relation, with values up to $10^{10} M_{\odot}$ (Lauer et al. 2007).

This discrepancy is investigated here by first looking at the upper extremity of the $M_{\text{bh}}-\sigma$ relation. At 400 km s^{-1} , it turns out that both the old $M_{\text{bh}}-\sigma$ relation — as given by Tremaine et al.’s (2002) regression of $\log M_{\text{bh}}$ on $\log \sigma$ — and the new relation (Eq.12) predict the same black hole mass: $2.3^{+1.7}_{-1.0} \times 10^9 M_{\odot}$. A value of 400 km s^{-1} is used here due to the rapid decline in the number density of systems with higher velocity dispersions (Sheth et al. 2003; Bernardi et al. 2006). This upper black hole mass agrees well with that from the $M_{\text{bh}}-n$ relation in GD07, where $M_{\text{bh,upper}} = 1.2^{+2.6}_{-0.4} \times 10^9 M_{\odot}$, implying an upper (1σ) SMBH mass limit of $\sim 4 \times 10^9 M_{\odot}$.

For the (K -band) $M_{\text{bh}}-L$ relation, $\log(M_{\text{bh}}/M_{\odot}) = (8.29 \pm 0.08) - (0.37 \pm 0.04)[M_K + 24]$ (Graham 2007), to predict a more massive black hole than $2.3 \times 10^9 M_{\odot}$ requires a spheroid with $M_K < -26.9$ mag. From the K -band magnitudes for 102 brightest cluster galaxies (BCGs) in Stott et al. (2007), while many galaxies are close to this limit, only two are brighter (after the small adjustment of 0.1 mag when switching from $H_0 = 70$ to 73 km s Mpc^{-1}). Furthermore, from the (corrected) SDSS r' -band BCG magnitudes in both Desroches et al. 2007, their Figure 9) and Liu et al. (2008, their Figure 13) we see that the brightest magnitudes truncate at $M_{r'} \sim -24.2$ mag. Using $r' - K = 2.8$, this corresponds to a K -band magnitude of -27.0 mag and an ($M_{\text{bh}}-L$)-derived SMBH mass of $2.5 \times 10^9 M_{\odot}$. It therefore appears that the $M_{\text{bh}}-L$ relation does not predict higher SMBH

masses than the $M_{\text{bh}}-\sigma$ relation. The near-infrared analysis by Batcheldor et al. (2007) also supports this picture (but see Lauer et al. 2007). The $M_{\text{bh}}-M_{\text{bh}}$ diagram from Lauer et al. (2007, their Fig.2) is at odds with the above result. It is however first noted that no non-BCG in Lauer et al. has a magnitude brighter than NGC 6876 at $M_V = -23.49$ mag ($H_0 = 73$). Assuming a $V - K$ color of 3.22 for elliptical galaxies (Buzzoni 2005), this magnitude corresponds to $M_K = -26.71$ mag, giving a black hole mass of $2.0 \times 10^9 M_\odot$, consistent with the upper bound from the $M_{\text{bh}}-\sigma$ relation.

To try and resolve the issue with the BCGs in Lauer et al., their $M_{\text{bh}}-M_{\text{bh}}$ diagram is reproduced here after applying a number of updates. First, the new $M_{\text{bh}}-\sigma$ relation (Eq.12) is applied to the velocity dispersions tabulated in Lauer et al.⁶. Second, the above mentioned K -band $M_{\text{bh}}-L$ relation is used (and $V - K = 3.22$ applied). As detailed in Graham (2007), this updated relation benefits from a number of factors, including (i) the identification of lenticular galaxies previously treated as elliptical galaxies, (ii) it was constructed in the near-infrared rather than the B - or V -band and so the magnitudes are less prone to biases from dust attenuation and young stellar populations, and (iii) a careful Sérsic bulge plus exponential disk decomposition has been performed. Graham (2007)⁷, report that a *symmetrical* regression of the M_{bh} and L_K data yields a slope of -0.40 . Given that the K -band stellar mass-to-light ratio is roughly constant for elliptical galaxies (Chabrier 2003; Bruzual & Charlot 2003), this corresponds to a one-to-one stellar-to-black hole mass relation — which makes sense at the high-mass end as it is the value expected from the dry merging of galaxies and the coalescence of their SMBHs. The V -band $M_{\text{bh}}-L$ relation used by Lauer et al. was much steeper than this, having a slope of -0.53 (i.e. $M_{\text{bh}} \propto L^{1.32}$), and subsequently predicted notably more massive SMBH masses. In addition, given that the V -band stellar mass-to-light ratio increases with luminosity, an even steeper dependence of the SMBH mass on stellar mass would be inferred, at odds with dry merging of massive galaxies. The results of applying the new $M_{\text{bh}}-L$ and $M_{\text{bh}}-\sigma$ relations to predict the SMBH masses for Lauer et al.’s data are shown in Fig.8a.

While Lauer et al. correctly used bulge magnitudes for the disc galaxies in their Fig.2, these were obtained from $R^{1/4}$ bulge plus exponential disc decompositions. Because most bulges have a Sérsic (1963) $R^{1/n}$ light profile (Graham & Driver 2005) with $n < 4$ (e.g., Graham 2001; Balcells et al. 2003; MacArthur et al. 2003), it is well known that such an approach overestimates the flux (e.g., Brown et al. 2003). To account for this, the bulge-to-disk flux ratios from Graham & Driver (2007b, their Table 2) have been applied⁸. This entailed reducing the S0 bulge magnitudes by $2.5 \log(0.25/0.60)$ and the Sa-Sb

bulge magnitudes by $2.5 \log(0.17/0.33)$. The results of doing this are shown in Fig.8b. This resolves the conflict seen in Fig.8a at the low-mass end. While such a correction is okay in a statistical sense, ideally individual galaxy corrections should be applied and this may well account for the increased scatter about the one-to-one line in Fig.8b.

Although the BCG tend to have ($M_{\text{bh}}-L$)-derived black hole masses smaller than $4 \times 10^9 M_\odot$, the Lauer et al. BCG magnitudes do tend to produce SMBH mass estimates that are roughly twice as large as those predicted from their velocity dispersions. From Liu et al.’s (2008) figure 5, one can see that the stellar envelope which surrounds (some) BCGs becomes significant (albeit relative to an $R^{1/4}$ model) at $\mu_{r'} \sim 23$ mag arcsec $^{-2}$, while Gonzalez et al. (2005, their Figure 3) indicates a value around 23.5 to 25 r' -mag arcsec $^{-2}$. These ranges are in agreement with the values seen in Seigar et al. 2007. This halo of stars is very likely, at least in part, due to stars that have been tidally stripped from galaxies within the cluster environment (e.g. Merritt et al. 1985). As such, it pertains more to the cluster than the BCG, and should be excluded from measurements of the BCG luminosity.

To avoid the issue of the outer envelope — which was thought to occur at $\mu_{r'} \sim 25$ mag arcsec $^{-2}$ — Lauer et al.’s BCG magnitudes were obtained from $R^{1/4}$ model fits to surface brightness profiles brighter than $\mu_{r'} = 23.74$ mag arcsec $^{-2}$ (Graham et al. 1996, using $r' - R_c = 0.24$). The danger is that some of the outermost portion of the light profile which was modeled may have been elevated to a brighter level by the flux of the envelope. As the light profiles did not extend to large radii, the ‘break’ in the profile, where the envelope starts to dominate, may have been missed. If so, such contamination would result in the best-fitting $R^{1/4}$ model having an increased effective radius and a brighter total flux. Therefore, before concluding a problem exists with the BCGs, it would be prudent to actually perform a galaxy/envelope decomposition (e.g. Gonzalez et al. 2005; Seigar et al. 2007) of such systems (not just those in Lauer et al.), enabling one to quantify how the envelope flux might be biasing the magnitudes. This is however beyond the scope of this paper.

It is perhaps worth noting that a few galaxies are known to possess both a SMBH and a nuclear star cluster, leading one to wonder whether a) the combined masses should be used and b) do the ‘offset’ barred galaxies have significant nuclear star clusters? As noted in GD07, the barred galaxy NGC 3384 has $M_{\text{NC}}/M_{\text{bh}} \sim 2$. Using $M_{\text{NC}} + M_{\text{bh}}$ rather than only M_{bh} would bring this galaxy back in line with the $M_{\text{bh}}-\sigma$ relation. However the barred galaxy NGC 1023 has an offset of nearly -0.7 dex but $M_{\text{NC}}/M_{\text{bh}} \sim 0.1$, while the barred galaxy NGC 2778 has no nuclear cluster but is offset by -0.9 dex. Furthermore, the unbarred galaxy NGC 7457, which has no offset from the $M_{\text{bh}}-\sigma$ relation has $M_{\text{NC}}/M_{\text{bh}} \sim 10$ (GD07). Nonetheless, a careful quantitative analysis of the nuclear structure of galaxies with SMBHs would be highly desirable; it would additionally enable one to explore the $M_{\text{bh}} - \text{central surface brightness relation}$ proposed by GD07.

At 100 km s^{-1} the barless $M_{\text{bh}}-\sigma$ relation yields masses (for unbarred galaxies) which are 67% higher than the expression in Tremaine et al. (2002). Past efforts to mea-

⁶ While this is not ideal because galaxies with bars may have their SMBH masses over-estimated, at the high mass end we do not predict larger SMBH masses than the $M_{\text{bh}}-\sigma$ relation from Tremaine et al. (2002) on this same data.

⁷ Kormendy & Gebhardt (2001) report a B -band $M_{\text{bh}}-L$ relation with a slope equal to -0.43 . Graham (2007) find a consistent slope of -0.38 ± 0.06 using the regression analysis from Tremaine et al. (2002).

⁸ An extensive analysis of dust-corrected bulge-to-disk systematics with disk galaxy type can be found in Graham & Worley (2008).

sure the SMBH mass function and mass density using the old $M_{\text{bh}}-\sigma$ relation (see the roundup in Graham & Driver 2007b, their table 3) may therefore need tweaking. Consequences for the $M_{\text{NC}}-\sigma$ (and $M_{\text{NC}}-L$) relation involving nuclear star clusters (e.g., Ferrarese et al. 2006; Balcells et al. 2007) are also deferred for elsewhere.

The Author hereby thanks Alessandro Marconi and

Leslie Hunt for kindly providing the major-to-minor axis ratios from their GALFIT bulge models. This research has made use of the NASA/IPAC Extragalactic Database (NED) and HyperLeda. This paper has benefited from two simultaneous referees, one of whom suggested that galaxies with partially edge-on bars may have their bulge R_e values underestimated.

REFERENCES

- Aguerri, J. A. L., Prieto, M., Varela, A. M., & Muñoz-Tuñón, C. 2001, *Ap&SS*, 276, 611
- Akritis, M.G., & Bershad, M.A., 1996, *ApJ*, 470, 706
- Aller, M.C., Richstone, D.O. 2007, *ApJ*, 665, 120
- Atkinson, J. W., et al. 2005, *MNRAS*, 359, 504
- Balcells, M., Graham, A.W., Dominguez-Palmero, L., Peletier, R.F., 2003, *ApJ*, 582, L79
- Balcells, M., Graham, A.W., Peletier, R.F., 2007, *ApJ*, 665, 1084
- Barway, S., Kembhavi, A. 2007, *ApJ*, 662, L67 (BK07)
- Batcheldor, D., Marconi, A., Merritt, D., & Axon, D. J. 2007, *ApJ*, 663, L85
- Bekki, K., Couch, W. J., Drinkwater, M. J., & Gregg, M. D. 2001, *ApJ*, 557, L39
- Bernardi, M., et al. 2006, *AJ*, 131, 2018
- Bernardi, M., Sheth, R.K., Tundo, E., Hyde, J.B., 2007, *ApJ*, 660, 267
- Blais-Ouellette, S., Amram, P., Carignan, C., & Swaters, R. 2004, *A&A*, 420, 147
- Brown, R.J.N., et al. 2003, *MNRAS*, 341, 747
- Bruzual, G., & Charlot, S. 2003, *MNRAS*, 344, 1000
- Busarello, G., Capaccioli, M., D'Onofrio, M., Longo, G., Richter, G., & Zaggia, S. 1996, *A&A*, 314, 32
- Buzzoni, A., 2005, *MNRAS*, 361, 725
- Caon, N., Capaccioli, M., D'Onofrio, M., 1993, *MNRAS*, 265, 1013
- Cappellari, M., & Emsellem, E. 2004, *PASP*, 116, 138
- Cappellari, M., Verolme, E.K., van der Marel, R.P., Verdoes Kleijn, G.A., Illingworth, G.D., Franx, M., Carollo, C.M., & de Zeeuw, P.T., 2002, *ApJ*, 578, 787
- Chabrier, G. 2003, *PASP*, 115, 763
- Coccatto, L., Sarzi, M., Pizzella, A., Corsini, E. M., Dalla Bontà, E., & Bertola, F. 2006, *MNRAS*, 366, 1050
- Cretton, N., & van den Bosch, F.C. 1999, *ApJ*, 514, 704
- Dalla Bontà, E., Ferrarese, L., Corsini, E. M., Miralda-Escude, J., Coccatto, L., & Pizzella, A. 2007, *Mem. S.A.It.*, 75, 282 (arXiv:0706.1959)
- Davies, R. I., et al. 2006, *ApJ*, 646, 754
- Debattista, V. P., Corsini, E. M., & Aguerri, J. A. L. 2002, *MNRAS*, 332, 65
- de Francesco, G., Capetti, A., & Marconi, A. 2006, *A&A*, 460, 439
- Desroches, L.-B., Quataert, E., Ma, C.-P., & West, A. A. 2007, *MNRAS*, 377, 402
- Djorgovski, G., & Davis, M. 1987
- D'Onofrio, M., Capaccioli, M., & Caon, N. 1994, *MNRAS*, 271, 523
- Dressler, A., Lynden-Bell, D., Burstein, D., Davies, R. L., Faber, S. M., Terlevich, R., & Wegner, G. 1987, *ApJ*, 313, 42
- Erwin, P., 2004, *A&A*, 415, 941
- Erwin, P., Beltrán, J. C. V., Graham, A. W., & Beckman, J. E. 2003, *ApJ*, 597, 929
- Eskridge, P.B., et al. 2000, *AJ*, 119, 536
- Faber, S. M., & Jackson, R. E. 1976, *ApJ*, 204, 668
- Falcke, H., Kording, E., Markoff, S., 2004, *A&A*, 414, 895
- Feigelson, E.D., Babu, G.J., 1992, *ApJ*, 397, 55
- Ferrari, F., Dottori, H., Caon, N., Nobrega, A., & Pavani, D. B. 2004, *MNRAS*, 347, 824
- Ferrarese, L., et al. 2006, *ApJ*, 644, L21
- Ferrarese, L., & Ford, H., 2005, *Space Science Reviews*, 116, 523 (FF05)
- Ferrarese, L., Merritt, D. 2000, *ApJ*, 539, L9
- Fukugita, M., Shimasaku, K., & Ichikawa, T. 1995, *PASA*, 107, 945
- Gebhardt, K., et al. 2000, *AJ*, 119, 1157
- Gebhardt, K., et al. 2003, *ApJ*, 583, 92
- Gonzalez A. H., Zabludoff A. I., Zaritsky D., 2005, *ApJ*, 618, 195
- Graham, A.W. 2001, *ApJ*, 121, 820
- Graham, A.W. 2002, *ApJ*, 568, L13
- Graham, A.W. 2007, *MNRAS*, 379, 711
- Graham, A.W., Driver, S.P. 2005, *PASA*, 22(2), 118 (astro-ph/0503176)
- Graham, A.W., Driver, S.P. 2007a, *ApJ*, 655, 77 (GD07)
- Graham, A.W., Driver, S.P. 2007b, *MNRAS*, 380, L15
- Graham, A.W., Driver, S.P., Allen, P.D., Liske, J. 2007, *MNRAS*, 378, 198
- Graham, A.W., Lauer, T.R., Colless, M.M., Postman, M. 1996, *ApJ*, 465, 534
- Greenhill, L. 2003, *ApJ*, 590, 162
- Greenhill, L.J., Moran, J.M., & Herrnstein, J.R. 1997, *ApJ*, 481, L23
- Häring, N., Rix, H.-W. 2004, *ApJ*, 604, L89
- Ho, L. C., Filippenko, A. V., & Sargent, W. L. W. 1997, *ApJ*, 487, 591
- Hopkins, P.F., Hernquist, L., Cox, T.J., Robertson, B., Krause, E., 2007, *ApJ*, in press (arXiv0707.4005)
- Houghton, R. C. W., Magorrian, J., Sarzi, M., Thatte, N., Davies, R. L., & Krajnović, D. 2006, *MNRAS*, 367, 2
- Huré, J.-M. 2002, *A&A*, 395, L21
- Jerjen, H., Binggeli, B., & Barazza, F.D. 2004, *AJ*, 127, 771
- Knapen, J. H., Shlosman, I., & Peletier, R. F. 2000, *ApJ*, 529, 93
- Kormendy, J. 1988, *ApJ*, 335, 40
- Kormendy, J., et al. 1997, *ApJ*, 482, L139
- Kormendy, J., Gebhardt, K., Fisher, D. B., Drory, N., Macchetto, F. D., & Sparks, W. B. 2005, *AJ*, 129, 2636
- Lanzoni, B., & Ciotti, L. 2003, *A&A*, 404, 819
- Lauer, T.R., et al. 2007, *ApJ*, 662, 808
- Liu, F. S., Xia, X. Y., Mao, S., Wu, H., & Deng, Z. G. 2008, *MNRAS*, submitted (arXiv:0712.0927)
- Lodato, G., & Bertin, G. 2003, *A&A*, 398, 517
- López-Corredoira, M., Cabrera-Lavers, A., Mahoney, T. J., Hammersley, P. L., Garzón, F., & González-Fernández, C. 2007, *AJ*, 133, 154
- MacArthur, L. A., Courteau, S., & Holtzman, J. A. 2003, *ApJ*, 582, 689
- Maciejewski, W., & Binney, J., 2001, *MNRAS*, 323, 831
- Marconi, A., Hunt, L.K. 2003, *ApJ*, 589, L21 (MH03)
- Marconi, A., et al. 2003, *ApJ*, 586, 868
- Marconi, A., et al. 2006, *A&A*, 448, 921
- Marinova, I., & Jogee, S. 2007, *ApJ*, 659, 1176
- Márquez, I., Masegosa, J., Durret, F., González Delgado, R. M., Moles, M., Maza, J., Pérez, E., & Roth, M. 2003, *A&A*, 409, 459
- McLure, R.J., Dunlop, J.S., 2002, *MNRAS*, 331, 795
- Merloni, A., Heinz, S., & di Matteo, T. 2003, *MNRAS*, 345, 1057
- Merritt, D. 1985, *ApJ*, 289, 18
- Merritt, D., & Ferrarese, L. 2001, *ApJ*, 547, 140
- Meusinger, H., & Ismail, H. A. 2007, *Astronomische Nachrichten*, 328, 562
- Mulchaey, J. S., & Regan, M. W. 1997, *ApJ*, 482, L135
- Nowak, N., Saglia, R. P., Thomas, J., Bender, R., Pannella, M., Gebhardt, K., & Davies, R. I. 2007, *MNRAS*, 379, 909
- Ohta, K., Aoki, K., Kawaguchi, T., & Kiuchi, G. 2007, *ApJS*, 169, 1
- Onken, C. A., et al. 2007, *ApJ*, 670, 105
- Pastorini, G., et al. 2007, *A&A*, 469, 405
- Peng, C. Y., Ho, L. C., Impey, C. D., & Rix, H.-W. 2002, *AJ*, 124, 266
- Pignatelli, E., Salucci, P., & Danese, L. 2001, *MNRAS*, 320, 124
- Rest, A., et al. 2001, 121, 2431
- Richstone et al. 2007, *ApJL*, submitted, astro-ph/0403257
- Rix, H.-W., Carollo, C.M., Freeman, K. 1999, *ApJ*, 513, L25
- Sarzi, M., et al. 2001, *ApJ*, 550, 65
- Seigar, M.S., Graham, A.W., Jerjen, H. 2007, *MNRAS*, 378, 1575
- Sérsic, J.-L., 1963, *Boletín de la Asociacion Argentina de Astronomia*, vol.6, p.41

- Shapiro, K. L., Cappellari, M., de Zeeuw, T., McDermid, R. M., Gebhardt, K., van den Bosch, R. C. E., & Statler, T. S. 2006, MNRAS, 370, 559
- Sheth, R.K., et al. 2003, ApJ, 594, 225
- Sil'chenko, O. K. 1999, AJ, 117, 2725
- Stott, J.P., Edge, A.C., Smith, G.P., Swinbank, A.M., & Ebeling, H. 2007, MNRAS, submitted (arXiv:0712.0496)
- Tonry, J., et al., 2001, ApJ, 546, 681
- Tremaine, S., et al., 2002, ApJ, 574, 740
- Trujillo, I., Erwin, P., Asensio Ramos, A., Graham, A.W. 2004, AJ, 127, 1917
- Trujillo, Graham, A.W., Caon, N. 2001, MNRAS, 326, 869
- Tundo, E., Bernardi, M., Hyde, J.B., Sheth, R.K., Pizzella, A., 2007, ApJ, 663, 53
- van Albada, G.D. 1980, 90, 123
- van der Marel, R.P., & van den Bosch, F.C. 1998, AJ, 116, 2220
- von der Linden, A., Best, P. N., Kauffmann, G., & White, S. D. M. 2007, MNRAS, 379, 867
- Virani, S. N., De Robertis, M. M., & VanDalsen, M. L. 2000, AJ, 120, 1739
- Wold, M., Lacy, M., Käuff, H. U., & Siebenmorgen, R. 2006, A&A, 460, 449
- Wozniak, H., Combes, F., Emsellem, E., & Friedli, D. 2003, A&A, 409, 469
- Wyse, R. F. G. 2004, ApJ, 612, L17
- Yu, Q., Tremaine, S., 2002, MNRAS, 335, 965

TABLE 1
RADII AND SURFACE BRIGHTNESSES

Galaxy	Type	b/a	R_e	$\langle\mu\rangle_{e,K}$
‘Group 1’				
NGC 4258	Sp,bar	0.51	0.66	15.26
NGC 4486 (M87)	E	0.87	6.0	16.86
NGC 3115	S0	0.38	2.9	16.49
NGC 4649	E	0.80	7.2	17.08
NGC 3031 (M81)	Sp	0.76	3.0	16.83
NGC 4374 (M84)	E	0.91	7.8	17.35
NGC 221 (M32)	S0(?)	0.71	(0.20)	(15.30)
NGC 5128 (Cen A)	S0	0.85	3.3	16.68
NGC 4697	E	0.58	6.9	18.18
IC 1459	E	0.75	7.1	16.95
NGC 5252	S0	0.48	6.7	17.19
NGC 2787	S0,bar	0.70	0.27	14.41
NGC 4594	Sp	0.63	4.0	16.21
NGC 3608	E	0.82	3.9	17.44
NGC 3245	S0	0.61	1.0	15.32
NGC 4291	E	0.77	2.0	16.22
NGC 3377	E	0.51	3.9	17.91
NGC 4473	E	0.57	2.1	16.41
Cygnus A	E	0.86	28.7	18.78
NGC 4261	E	0.80	5.8	16.82
NGC 4564	S0	0.66	(2.4)	(17.12)
NGC 4742	E	0.63	1.6	16.59
NGC 3379	E	0.87	2.7	16.54
NGC 1023	S0,bar	0.80	1.1	15.23
NGC 5845	E	0.70	0.42	13.70
NGC 3384	S0,bar	0.94	0.48	14.36
NGC 6251	E	0.82	10.0	17.06
‘Group 2’				
Milky Way	Sp,bar	1.00	0.7	15.54
NGC 224 (M31)	Sp	1.00	1.0	15.77
NGC 1068	Sp	0.89	2.9	15.91
NGC 4459	S0	0.83	13.7	19.76
NGC 4596	S0,bar	0.87	1.5	15.66
NGC 7457	S0	0.64	3.8	19.70
NGC 4342	S0	0.52	0.21	14.48
NGC 821	E	0.62	15.7	19.78
NGC 2778	S0	0.77	(2.6)	(17.69)
NGC 7052	E	0.57	9.1	17.32

NOTE. — Column 3: Spheroidal component’s minor-to-major axis ratio, b/a , for the galaxies from Marconi & Hunt (2003, priv. comm.). Columns 4 and 5: Geometric mean radii ($R_e = \sqrt{R_{e,maj}^2(b/a)}$) in kpc, and mean K -band effective surface brightnesses (Eq.3) in mag arcsec $^{-2}$, for the spheroidal component. A bracketed entry reflects that no (K -band) bulge/disk decomposition was performed.

TABLE 2
NEW AND UPDATED SMBH DATA

Galaxy	Type	Dist. (Mpc)	σ km s ⁻¹	log R_e [kpc]	M_{bh} ($10^8 M_\odot$)
1	2	3	4	5	6
Updated Marconi & Hunt (2003) data					
Cygnus A	E	240	270	1.49	$26.0^{+7.0}_{-7.0}$ [2]
Cen A	S0	4.2	150	0.56	$1.1^{+0.1}_{-0.1}$ [10]
NGC 221	S0	0.8	75	-0.98 [6]	$0.025^{+0.005}_{-0.005}$
NGC 821	E	24.1	200 [5]	1.30	$0.85^{+0.35}_{-0.35}$ [11]
NGC 2778	S0	22.9	175	-0.60 [2]	$0.14^{+0.08}_{-0.09}$
NGC 3379	E	10.6	206	0.46	$1.4^{+2.6}_{-1.0}$ [12]
NGC 4342	S0	17.0 [1]	251 [5]	-0.36	$3.3^{+1.9}_{-1.1}$
NGC 4374	E	18.4	296	0.91	$4.64^{+3.46}_{-1.83}$ [13]
NGC 4564	S0	15.0	162	-0.50 [2]	$0.56^{+0.03}_{-0.08}$
NGC 5252	S0	94.4 [2]	190	0.98	$9.7^{+14.9}_{-4.6}$
NGC 6251	E	101 [2]	290	1.02	$5.8^{+1.8}_{-2.0}$
NGC 7052	E	60.0 [2]	266	1.00	$3.4^{+2.4}_{-1.3}$
New galaxies with SMBH measurements					
NGC 1300	SBbc	$19.3h_{73}$	229 [5]	-0.28 [7]	$0.68^{+0.65}_{-0.33}$ [14]
NGC 1399	E	20.0 [3]	317	1.09 [4]	12^{+5}_{-6} [15]
NGC 2748	Sbc	$23.8h_{73}$	...	...	$0.45^{+0.36}_{-0.37}$ [14]
NGC 3227	SB	$17.0h_{73}$	160	-0.57	$0.15^{+0.05}_{-0.08}$ [16]
NGC 3998	S0	14.1 [3]	305	-0.16	$2.2^{+2.0}_{-1.7}$ [17]
NGC 4151	SBab	13.9	156 [5]	-0.23 [8]	$0.45^{+0.05}_{-0.05}$ [18]
NGC 4435	SB0	16.0	157	-0.07	< 0.075 [19]
NGC 4486a	E	17.0 [1]	110	-0.39 [9]	$0.13^{+0.08}_{-0.08}$ [20]
NGC 7582	SBab	22.4	156	...	$0.55^{+0.26}_{-0.19}$ [21]

NOTE. — Unless otherwise specified, the distances, velocity dispersions, and effective radii of each galaxy have come from the reference which provides the SMBH mass. Both M_{bh} and R_e have been adjusted to the distance given in column 2. References: 1 = Jerjen et al. (2004); 2 = Graham & Driver (2007a); 3 = Tonry et al. (2001); 4 = D’Onofrio et al. (1994); 5 = HyperLeda (<http://leda.univ-lyon1.fr/>); 6 = Graham (2002); 7 = Aguerri et al. (2001); 8 = Virani et al. (2000); 9 = Kormendy et al. (2005); 10 = Marconi et al. (2006); 11 = Richstone et al. (2007); 12 = Shapiro et al. (2006); 13 = Maciejewski & Binney (2001); 14 = Atkinson et al. (2005); 15 = Houghton et al. (2006); 16 = Davies et al. (2006); 17 = De Francesco et al. (2006); 18 = Onken et al. (2007); 19 = Coccato et al. (2006); 20 = Nowak et al. (2007); 21 = Wold et al. (2006).

TABLE 3
GALAXIES WITH SOMEWHAT UNCERTAIN M_{bh}

Galaxy	Reference	Note
Abell 1836	Dalla Bontà et al. 2007	no refereed publication
Abell 3565	Dalla Bontà et al. 2007	no refereed publication
Circinus	Greenhill et al. 2003	poorly known disk inclination
IC 1459	Cappellari et al. 2002	gas/stellar dynamics differ
NGC 1068	Huré 2002; Lodato & Bertin 2003	disk model uncertain
NGC 4041	Marconi et al. 2003	disk might be dynamically decoupled
NGC 4303	Pastorini et al. 2007	poorly known disk inclination
NGC 4350	Pignatelli et al. 2001	possible SMBH, high $M_{\text{bh}}/M_{\text{bulge}}$
NGC 4459	Sarzi et al. 2001	poorly known disk inclination
NGC 4486B	Kormendy et al. 1997	possible SMBH, $M_{\text{bh}}/M_{\text{bulge}} = 0.09$
NGC 4594	Kormendy 1988	no 3-integral model
NGC 4596	Sarzi et al. 2001	poorly known disk inclination
NGC 4945	Greenhill et al. 1997	no 2D velocity field
NGC 5055	Blais-Ouellette et al. 2004	possibly no black hole
NGC 7332	Håring & Rix 2004	no refereed publication

TABLE 4
SMBH MASS – SPHEROID RELATIONS

Sample	Relation	Δ dex	Δ' dex
$M_{\text{bh}}-\sigma$			
40 galaxies	$8.13 \pm 0.06 + (3.92 \pm 0.27) \log(\sigma/200)$	0.38	0.35
29 non-barred	$8.26 \pm 0.06 + (3.67 \pm 0.19) \log(\sigma/200)$	0.30	0.25
19 elliptical	$8.25 \pm 0.05 + (3.68 \pm 0.25) \log(\sigma/200)$	0.24	0.18
$M_{\text{bh}}-\sigma-R_e$			
40 galaxies	$8.19 \pm 0.05 + (3.23 \pm 0.28) \log(\sigma/200) + (0.43 \pm 0.11) [\log(R_e/3)]$	0.30	0.28
29 non-barred	$8.26 \pm 0.05 + (3.29 \pm 0.26) \log(\sigma/200) + (0.29 \pm 0.11) [\log(R_e/3)]$	0.25	0.24
19 elliptical	$8.23 \pm 0.04 + (3.32 \pm 0.36) \log(\sigma/200) + (0.21 \pm 0.15) [\log(R_e/3)]$	0.22	0.18

NOTE. — The total scatter Δ is given rather than the (smaller) internal/intrinsic scatter, as the latter quantity depends on the measurements errors that one assigns. The final column shows the total scatter Δ' after removing just two data points (Cygnus A and NGC 3998).




FIG. 1.— Using the 27 galaxies from GD07, the residuals about their $M_{\text{bh}}\text{-}\sigma$ relation (constructed to minimize the scatter in the $\log M_{\text{bh}}$ direction) are shown against their Sérsic index n . Panel a) highlights the 5 galaxies with an unresolved SMBH sphere-of-influence (Table II in FF05) — they are not responsible for the trend. Panel b) shows the 13 disk galaxies (circled); they are responsible for the trend seen in this figure.




FIG. 2.— Similar to Fig. 1 except that the $\delta \log M_{\text{bh}}$ values are shown against a) the geometric mean effective radii from Table 1, b) the Sérsic indices n from GD07, and c) the absolute K -band magnitudes M_K from MH03. The barred galaxies have been circled; they are responsible for the bulk of the apparent trends.




FIG. 3.— Panel a) $M_{\text{bh}}-L$ relation for the 27 galaxies from GD07. Panel b) shows the residuals versus their geometric mean effective radii. The six barred galaxies have been circled.




FIG. 4.— Panel a) Residuals about the $M_{\text{bh}}-\sigma$ relation using the 27 ‘Group 1’ galaxies from MH03 plus their data for the Milky Way and M31 are shown against their tabulated major-axis effective radii. The five barred galaxies are denoted with crosses. Panel b) Residuals about the $M_{\text{bh}}-\sigma$ relation when using only the elliptical galaxies (according to MH03) to construct the $M_{\text{bh}}-\sigma$ relation. The non-elliptical galaxies NGC 221 and NGC 4564 have been circled. The point in the top right is Cygnus A at 240 Mpc (using $M_{\text{bh}} = 2.9 \times 10^9 M_{\odot}$).




FIG. 5.— $M_{\text{bh}}\text{-}\sigma$ diagram for 40 galaxies (see section 4.4). The 11 barred galaxies are denoted with a cross.




FIG. 6.— Panel a) Residuals about the $M_{\text{bh}}\text{-}\sigma$ relation constructed using the 29 non-barred galaxies (see section 4.4). The residual offset of the ten barred galaxies that have R_e values are denoted with a cross. Panel b) Residuals about the $M_{\text{bh}}\text{-}\sigma$ relation constructed using the 19 elliptical galaxies.




FIG. 7.— The R_c -band σ - L relation for SDSS early-type galaxies (dashed line) taken from Tundo et al. (2007, their Eq.4), and for non-barred galaxies with direct SMBH mass estimates in MH03 (solid line, Eq.13). The outlying galaxy NGC 4342 has been excluded.




FIG. 8.— M_{bh} masses for the galaxies tabulated in Lauer et al. (2007) obtained using the new $M_{\text{bh}}-\sigma$ relation ((Eq.12) and the $M_{\text{bh}}-L$ relation from Graham (2007, his Eq.14). In panel b) the $R^{1/4}$ bulge magnitudes of the disc galaxies have been adjusted as described in Section ?? . The BCGs and normal galaxies are denoted with open circles and dots, respectively.